

The Journal of Physiology

Ca²⁺ signalling in urethral interstitial cells of Cajal

Gerard P. Sergeant, M. A. Hollywood, N. G. McHale and K. D. Thornbury

J. Physiol. 2006;576;715-720; originally published online Aug 17, 2006;

DOI: 10.1113/jphysiol.2006.115956

This information is current as of November 13, 2007

This is the final published version of this article; it is available at:

<http://jp.physoc.org/cgi/content/full/576/3/715>

This version of the article may not be posted on a public website for 12 months after publication unless article is open access.

The Journal of Physiology Online is the official journal of The Physiological Society. It has been published continuously since 1878. To subscribe to *The Journal of Physiology Online* go to: <http://jp.physoc.org/subscriptions/>. *The Journal of Physiology Online* articles are free 12 months after publication. No part of this article may be reproduced without the permission of Blackwell Publishing: JournalsRights@oxon.blackwellpublishing.com

SYMPOSIUM REPORT

Ca²⁺ signalling in urethral interstitial cells of Cajal

Gerard P. Sergeant, M. A. Hollywood, N. G. McHale and K. D. Thornbury

Smooth Muscle Research Centre, Dundalk Institute of Technology, Dundalk, County Louth, Ireland

Interstitial cells of Cajal (ICC) in the urethra have been proposed as specialized pacemakers that are involved in the generation of urethral tone and therefore the maintenance of urinary continence. Recent studies on freshly dispersed ICC from the urethra of rabbits have demonstrated that pacemaker activity in urethra ICC is characterized by spontaneous transient depolarizations (STDs) under current clamp and spontaneous transient inward currents (STICs) under voltage clamp. When these events were simultaneously recorded with changes in intracellular Ca²⁺ (using a Nipkow spinning disk confocal microscope) they were found to be associated with global Ca²⁺ oscillations. In this short review we will consider some of these recent findings regarding the contribution of intracellular Ca²⁺ stores and Ca²⁺ influx to the generation of pacemaker activity in urethral ICC with particular emphasis on the contribution of reverse Na⁺/Ca²⁺ exchange (NCX).

(Received 26 June 2006; accepted after revision 11 August 2006; first published online 17 August 2006)

Corresponding author G. P Sergeant: Smooth Muscle Research Centre, Regional Development Centre, Dundalk Institute of Technology, Dundalk, Co. Louth, Ireland. Email: gerard.sergeant@dkit.ie

It is now well established that ICC in specific regions of the GI tract act as pacemakers that are responsible for generating electrical slow waves which form the basis of co-ordinated waves of contraction observed throughout the gut (Thuneberg, 1982; Sanders, 1996; Dickens *et al.* 1999; Sanders *et al.* 2006). Therefore, when ICC with similar morphological and immunohistochemical properties to those in the gut were found in the smooth muscle layers of the urethra, it was suggested that they may perform a similar function (Sergeant *et al.* 2000). This idea was strengthened with the results of electrophysiological experiments which showed that urethral ICC were spontaneously active and that the pattern of electrical activity in single ICC resembled that recorded from the strips of whole tissue using intracellular microelectrodes (Hashitani *et al.* 1996; Hashitani & Edwards, 1999; Sergeant *et al.* 2006). In contrast, the majority of smooth muscle cells (SMCs) isolated from the urethra were electrically quiescent and were therefore not considered as the source of the electrical activity recorded from the whole tissue. ICC in the urethra were originally referred to as interstitial cells (IC; Sergeant *et al.* 2000). However, in order to try to standardize the

terminology in the literature they are now referred to as ICC (Sergeant *et al.* 2006; Bradley *et al.* 2006) in line with those in the gastrointestinal (GI) tract.

Pacemaker mechanism

STICs in urethral ICC were inhibited by the chloride channel blockers A-9-C and niflumic acid (Sergeant *et al.* 2000), and their reversal potential closely followed the predicted chloride equilibrium potential (E_{Cl} , Sergeant *et al.* 2000, 2006) suggesting that they were due to activation of Ca²⁺-activated Cl⁻ channels. Although STICs with a similar pharmacological profile have been recorded from various smooth muscles, including rabbit portal vein and pulmonary artery (Wang *et al.* 1992; Hogg *et al.* 1993) and canine and guinea pig trachea (Janssen & Sims, 1994), the amplitude and temporal profile of the STICs recorded from urethra ICC were quite different. For example, STICs in isolated urethral ICC often exceeded 900 pA (Sergeant *et al.* 2001) compared to amplitudes of ~100 pA in SMCs (Large & Wang, 1996). In addition, the kinetics of STICs in ICC were much slower (> 1 s duration) in urethra ICC compared to their smooth muscle counterparts (< 100 ms duration). These differences were thought to reflect differences in the underlying Ca²⁺ signals responsible for generating STICs. Previous studies had indicated that STICs in SMCs were caused by Ca²⁺ sparks (ZhuGe *et al.* 1998; Gordienko *et al.* 1999). However, given the small single channel conductance of

This report was presented at The Journal of Physiology Symposium on Involvement of interstitial cells of Cajal in the control of smooth muscle excitability, Okayama, Japan, 22 July 2006. It was commissioned by the Editorial Board and reflects the views of the authors.

Ca^{2+} -activated Cl^- channels (2.6 pS; Klockner, 1993) it seemed unlikely that a Ca^{2+} spark would be able to activate a sufficient number of channels to generate STICs of the large amplitudes recorded in urethral ICC. Instead, it was proposed that STICs in ICC arose from global Ca^{2+} oscillations (Sergeant *et al.* 2001).

Role of Ca^{2+} stores

The importance of Ca^{2+} stores to pacemaker activity in urethra ICC was firmly established in a study by Sergeant *et al.* (2001) which demonstrated that the sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase (SERCA) inhibitor, cyclopiazonic acid (CPA) abolished STICs. Furthermore, activation of STICs appeared to involve release of Ca^{2+} from both inositol trisphosphate

(IP_3)- and ryanodine-sensitive stores as they were abolished by the phospholipase C (PLC) inhibitor 2-nitro-4-carboxyphenyl-*N,N*-diphenylcarbamate (NCDC), the IP_3 receptor (IP_3R) blocker, 2-aminoethoxydiphenyl borate (2-APB) (Maruyama *et al.* 1997) as well as caffeine and ryanodine. The contribution of IP_3Rs and ryanodine receptors (RyRs) to pacemaker activity in urethral ICC was assessed in more detail in a later study by Johnston *et al.* (2005) which looked at the effects of these agents on the Ca^{2+} oscillations underlying STICs. This study showed that urethra ICC loaded with fluo-4 AM developed regularly occurring global Ca^{2+} oscillations that were associated with STICs as originally predicted by Sergeant *et al.* (2001). Interestingly, while application of the RyR inhibitor tetracaine completely abolished these events, inhibition of IP_3Rs with 2-APB only decreased their spatial spread and converted the propagating Ca^{2+} waves into




Figure 1. Pseudo linescan or x,t plot of propagating Ca^{2+} waves in an isolated urethra ICC Ca^{2+} waves were abolished by removal of Ca_o^{2+} (A). Reduction in $[\text{Ca}^{2+}]_o$ from 1.8 to 0.9 mM reduced the frequency of Ca^{2+} waves by $\sim 50\%$ (B). An increase in $[\text{Ca}^{2+}]_o$ to 3.6 mM increased the frequency of Ca^{2+} waves (C). Modified from Sergeant *et al.* (2006).




Figure 2. Simultaneous recording of membrane potential in current clamp and $[Ca^{2+}]_i$ in a fluo-4 AM loaded urethral ICC, imaged with a Nipkow spinning disk confocal microscope shows that STDs (red) are associated with spontaneous Ca^{2+} oscillations (blue)

In the presence of 10 μM nifedipine the duration of both events is shortened, but their frequency is not affected.

more localized events. Therefore, it appeared that IP_3Rs were critically involved in propagation of Ca^{2+} waves, but that RyRs were responsible for generating the primary pacemaker event. The RyRs were therefore considered the 'prime oscillators' in the pacemaker mechanism.

Ca^{2+} influx in urethra ICC

Johnston *et al.* (2005) also demonstrated that the frequency of Ca^{2+} oscillations in urethra ICC was not only dependent on Ca^{2+} release from stores but was also critically dependent on Ca^{2+} entry. For example, when $[Ca^{2+}]_o$ was elevated to 3.6 mM the frequency of the Ca^{2+} waves increased significantly. Conversely, a reduction of $[Ca^{2+}]_o$ from 1.8 mM to 0.9 mM decreased wave frequency by ~40% and removal of $[Ca^{2+}]_o$ led to the immediate cessation of oscillations. An example of these effects is shown in Fig. 1. One possible interpretation of these results is that the reduction in $[Ca^{2+}]_o$ decreased the Ca^{2+} content of intracellular stores sufficiently to reduce wave frequency. However, Johnston *et al.* (2005) suggested that this was not the case. They showed that application of 10 mM caffeine for 10 s to a spontaneously active ICC evoked a large Ca^{2+} transient. When repeated in the presence Ca^{2+} free medium the amplitude of the caffeine response was unaffected despite the fact that spontaneous Ca^{2+} oscillations had ceased. This suggested that during the exposure to Ca^{2+} free medium (60 s) the intracellular Ca^{2+} stores remained intact.

This study also showed that the frequency of Ca^{2+} waves was inhibited by high concentrations (1 mM) of the non-specific Ca^{2+} entry blockers Cd^{2+} and La^{3+} but little affected by nifedipine (10 μM). More recent experiments

have examined the effect of nifedipine in more detail. Figure 2 shows a simultaneous recording of membrane potential in current clamp and $[Ca^{2+}]_i$ in a fluo-4 AM loaded urethral ICC, imaged with a Nipkow spinning disk confocal microscope. Under control conditions the cell exhibited regularly occurring STDs (shown in red) of ~70 mV amplitude comprising a spike and plateau at 0 mV. These events were associated with spontaneous Ca^{2+} oscillations (shown in blue). Application of nifedipine (10 μM) slightly decreased the amplitude and duration of STDs and Ca^{2+} oscillations, but their frequency was not affected suggesting that L-type Ca^{2+} channels are not involved in the generation of pacemaker activity.

Given the importance of Ca^{2+} stores to the generation of pacemaker activity, studies were also completed to assess the contribution of capacitative Ca^{2+} entry (CCE) to the influx pathway. Bradley *et al.* (2005) showed that isolated ICC exhibited a rise in $[Ca^{2+}]_i$ in response to addition of




Figure 3. Representative recording from an isolated ICC showing that reduction of $[Na^+]_o$ from 130 to 13 mM doubles the frequency of spontaneous Ca^{2+} oscillations

Modified from Bradley *et al.* (2006).




Figure 4. Representative recordings showing the effect of the selective reverse mode NCX inhibitors KB-R7943 and SEA0400 on spontaneous Ca²⁺ oscillations (A and B, respectively) and STICs recorded at -60 mV (C and D, respectively)

Modified from Bradley *et al.* (2006).

Ca²⁺ to the bath which was much larger when Ca²⁺ stores were depleted with CPA than under control conditions. This 'overshoot' in Ca²⁺ is a hallmark of CCE in a range of cell types (Hallam *et al.* 1989; Holda *et al.* 1998; Wilson

et al. 2002) indicating that urethral ICC possessed a similar CCE pathway. CCE in urethral ICC was inhibited by Ni²⁺ (100 μM), La³⁺ (10 μM) and Gd³⁺ (10 μM), but was not affected by SKF96365 (10 μM), nifedipine (10 μM) or




Figure 5. Schematic model of pacemaker activity in isolated urethra ICC

Ca²⁺ release from RyRs (red) is initiated by Ca²⁺ influx via reverse NCX. This in turn causes further release of Ca²⁺ from IP₃Rs (green) which are therefore responsible for propagation of these events. The raised intracellular [Ca²⁺] leads to stimulation of plasmalemmal Ca²⁺-activated Cl⁻ channels causing depolarization of the membrane and activation of L-type Ca²⁺ channels.

wortmanin ($1 \mu\text{M}$), and only slightly diminished (21%) by 2-APB ($100 \mu\text{M}$). Interestingly, the agents which inhibited CCE did not abolish STICs, suggesting that another Ca^{2+} entry pathway must be involved in the generation of these events.

A clue to the identity of a Ca^{2+} entry pathway involved came from an observation by Putney *et al.* (2001) who noted that in cells which don't display CCE, refilling of Ca^{2+} stores is most likely achieved by Ca^{2+} influx via reverse NCX. NCX is also known to be inhibited by high concentrations of La^{3+} and Cd^{2+} (Blaustein & Lederer, 1999) which had been previously shown to inhibit Ca^{2+} oscillations in urethra ICC (Johnston *et al.* 2005). This prompted us to investigate if Ca^{2+} influx via reverse NCX contributed to pacemaker activity in isolated urethral ICC. Although NCX is typically thought of as a Ca^{2+} extrusion mechanism it is in fact a bidirectional ion transport protein which can mediate Ca^{2+} entry depending on the net electrochemical driving force acting on it (Blaustein & Lederer, 1999). Bradley *et al.* (2006) tested if interventions designed to enhance reverse mode NCX increased the frequency of pacemaker activity. They found that reduction of $[\text{Na}^+]_o$ from 130 to 13 mM (which will make E_{NCX} more negative and therefore increase the driving force for Ca^{2+} entry via reverse NCX) dramatically increased the frequency of Ca^{2+} oscillations and STICs. A representative example of the effect of 13 mM $[\text{Na}^+]_o$ on spontaneous Ca^{2+} oscillations is shown in Fig. 3. Bradley *et al.* (2006) also reported that two putative reverse NCX inhibitors (KB-R7943 and SEA0400; Iwamoto *et al.* 1996; Watano *et al.* 1996; Matsuda *et al.* 2001; Lee *et al.* 2004) inhibited or else greatly reduced the frequency of both STICs at -60 mV and spontaneous Ca^{2+} oscillations. The latter effect was consistently accompanied by a fall in basal Ca^{2+} levels suggesting that Ca^{2+} influx via this pathway was involved in setting basal Ca^{2+} levels in these cells. Representative examples of these effects are shown in Fig. 4. These effects were noted to be similar to those caused by tetracaine, but not by 2-APB (Johnston *et al.* 2005).

Wu & Fry (2001) demonstrated that caffeine-induced Ca^{2+} release could be enhanced by Ca^{2+} influx via reverse NCX in guinea pig detrusor smooth muscle cells. However, Bradley *et al.* (2006) found that KB-R7943 and SEA0400 did not significantly affect Ca^{2+} -activated Cl^- currents (I_{ClCa}) evoked by application of caffeine or noradrenaline, suggesting that reverse NCX did not affect refilling of intracellular Ca^{2+} stores in urethra ICC. These data also demonstrated that KB-R7943 and SEA0400 did not produce their inhibitory effects via non-specific actions on Ca^{2+} -activated Cl^- channels, IP_3 Rs or RyRs. The effects of KB-R7943 and SEA0400 were notably similar to those produced by tetracaine (Johnston *et al.* 2005). However, as KB-R7943 and SEA0400 were shown not to inhibit RyRs directly, it suggested that Ca^{2+} influx via reverse NCX

triggered release of Ca^{2+} from RyRs. Such a mechanism was reminiscent of that described by Leblanc & Hume (1990) in cardiac myocytes. In these cells reverse mode NCX was facilitated by raised $[\text{Na}^+]_i$ due to influx via a TTX sensitive Na^+ channel. Thus far we have no evidence that such a pathway exists in urethral ICC; however, any mechanism that would raise $[\text{Na}^+]_i$ in the vicinity of the exchanger would tend to favour reverse mode NCX by making E_{NCX} more negative.

Model of pacemaker activity

Based on the currently available data one cycle of pacemaker activity can be considered to comprise the following sequence of events: (1) Ca^{2+} influx via reverse NCX, (2) activation of RyRs, causing localized Ca^{2+} release events, (3) propagation of these events by opening of IP_3 Rs, (4) stimulation of plasmalemmal Ca^{2+} -activated Cl^- channels causing depolarization, and (5) activation of L-type Ca^{2+} channels. This model is illustrated in the schematic diagram shown in Fig. 5.

Summary

ICC in the urethra have been proposed as pacemaker cells which may regulate spontaneous myogenic tone. In this short review we have attempted to assess some of the recent studies which have described different components which contribute to the pacemaker mechanism in isolated ICC. Studies in this field are at an early stage in comparison with those in the GI tract, but a clearer picture of the mechanisms involved is beginning to emerge and points to differences in the cellular basis of pacemaking in ICC in both regions. Further studies will be required to test the model of pacemaking proposed above, as well as to investigate the function of these cells in intact syncytia.

References

- Blaustein MP & Lederer WJ (1999). Sodium/calcium exchange: its physiological implications. *Physiol Rev* **79**, 763–854. (Review)
- Bradley E, Hollywood MA, Johnston L, Large RJ, Matsuda T, Baba A, McHale NG, Thornbury KD & Sergeant GP (2006). Contribution of reverse $\text{Na}^+/\text{Ca}^{2+}$ exchange to spontaneous activity in interstitial cells of Cajal in the rabbit urethra. *J Physiol* **574**, 651–661.
- Bradley E, Hollywood MA, McHale NG, Thornbury KD & Sergeant GP (2005). Pacemaker activity in urethral interstitial cells is not dependent on capacitative calcium entry. *Am J Physiol Cell Physiol* **289**, C625–C632.
- Dickens EJ, Hirst GD & Tomita T (1999). Identification of rhythmically active cells in guinea-pig stomach. *J Physiol* **514**, 515–531.
- Gordienko DV, Zholos AV & Bolton TB (1999). Membrane ion channels as physiological targets for local Ca^{2+} signalling. *J Microsc* **196**, 305–316.

- Hallam TJ, Jacob R & Merritt JE (1989). Influx of bivalent cations can be independent of receptor stimulation in human endothelial cells. *Biochem J* **259**, 125–129.
- Hashitani H & Edwards FR (1999). Spontaneous and neurally activated depolarizations in smooth muscle cells of the guinea-pig urethra. *J Physiol* **514**, 459–470.
- Hashitani H, Van Helden DF & Suzuki H (1996). Properties of spontaneous depolarizations in circular smooth muscle cells of rabbit urethra. *Br J Pharmacol* **118**, 1627–1632.
- Hogg RC, Wang Q, Helliwell RM & Large WA (1993). Properties of spontaneous inward currents in rabbit pulmonary artery smooth muscle cells. *Pflugers Arch* **425**, 233–240.
- Holda JR, Klishin A, Sedova M, Huser J & Blatter LA (1998). Capacitative calcium entry. *News Physiol Sci* **13**, 157–163.
- Iwamoto T, Watano T & Shigekawa M (1996). A novel isothiourea derivative selectively inhibits the reverse mode of $\text{Na}^+/\text{Ca}^{2+}$ exchange in cells expressing NCX1. *J Biol Chem* **271**, 22391–22397.
- Janssen LJ & Simms SM (1994). Spontaneous transient inward current and rhythmicity in canine and guinea-pig tracheal smooth muscle cells. *Pflugers Arch* **427**, 473–480.
- Johnston L, Sergeant GP, Hollywood MA, Thornbury KD & McHale NG (2005). Calcium oscillations in interstitial cells of the rabbit urethra. *J Physiol* **565**, 449–461.
- Klockner U (1993). Intracellular calcium ions activate a low-conductance chloride channel in smooth-muscle cells isolated from human mesenteric artery. *Pflugers Arch* **424**, 231–237.
- Large WA & Wang Q (1996). Characteristics and physiological role of the Ca^{2+} -activated Cl^- conductance in smooth muscle. *Am J Physiol* **271**, C435–454. (Review)
- Leblanc N & Hume JR (1990). Sodium current-induced release of calcium from cardiac sarcoplasmic reticulum. *Science* **248**, 372–376.
- Lee C, Visen NS, Dhalla NS, Le HD, Isaac M, Choptiany P, Gross G, Omelchenko A, Matsuda T, Baba A, Takahashi K, Hnatowich M & Hryshko LV (2004). Inhibitory profile of SEA0400 [2-[4-[(2,5-difluorophenyl) methoxy]phenoxy]-5-ethoxyaniline] assessed on the cardiac $\text{Na}^+/\text{Ca}^{2+}$ exchanger, NCX1.1. *J Pharmacol Exp Ther* **311**, 748–757.
- Maruyama T, Kanaji T, Nakade S, Kanno T & Mikoshiba K (1997). 2-Aminoethoxydiphenyl borate, a membrane-penetrable modulator of $\text{Ins}(1,4,5)\text{P}_3$ -induced Ca^{2+} release. *J Biochem* **122**, 498–505.
- Matsuda T, Arakawa N, Takuma K, Kishida Y, Kawasaki Y, Sakaue M, Takahashi K, Takahashi T, Suzuki T, Ota T, Hamano-Takahashi A, Onishi M, Tanaka Y, Kameo K & Baba A (2001). SEA0400, a novel and selective inhibitor of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger, attenuates reperfusion injury in the in vitro and in vivo cerebral ischemic models. *J Pharmacol Exp Ther* **298**, 249–256.
- Putney JW Jr (2001). Pharmacology of capacitative calcium entry. *Mol Interv* **1**, 84–94.
- Sanders KM (1996). A case for interstitial cells of Cajal as pacemakers and mediators of neurotransmission in the gastrointestinal tract. *Gastroenterology* **111**, 492–515.
- Sanders KM, Koh SD & Ward SM (2006). Interstitial cells of Cajal as pacemakers in the gastrointestinal tract. *Annu Rev Physiol* **68**, 307–343.
- Sergeant GP, Hollywood MA, McCloskey KD, McHale NG & Thornbury KD (2001). Role of IP_3 in modulation of spontaneous activity in pacemaker cells of rabbit urethra. *Am J Physiol Cell Physiol* **280**, C1349–C1356.
- Sergeant GP, Hollywood MA, McCloskey KD, Thornbury KD & McHale NG (2000). Specialised pacemaking cells in the rabbit urethra. *J Physiol* **526**, 359–366.
- Sergeant GP, Thornbury KD, McHale NG & Hollywood MA (2006). Interstitial cells of Cajal in the urethra. *J Cell Mol Med* **10**, 280–291.
- Thuneberg L (1982). Interstitial cells of Cajal: intestinal pacemaker cells? *Adv Anat Embryol Cell Biol* **71**, 1–130.
- Wang Q, Hogg RC & Large WA (1992). Properties of spontaneous inward currents recorded in smooth muscle cells isolated from the rabbit portal vein. *J Physiol* **451**, 525–537.
- Watano T, Kimura J, Morita T & Nakanishi H (1996). A novel antagonist, 7943, of the $\text{Na}^+/\text{Ca}^{2+}$ exchange current in guinea-pig cardiac ventricular cells. *Br J Pharmacol* **119**, 555–563.
- Wilson SM, Mason HS, Smith GD, Nicholson N, Johnston L, Janiak R & Hume JR (2002). Comparative capacitative calcium entry mechanisms in canine pulmonary and renal arterial smooth muscle cells. *J Physiol* **543**, 917–931.
- Wu C & Fry CH (2001). $\text{Na}^+/\text{Ca}^{2+}$ exchange and its role in intracellular Ca^{2+} regulation in guinea pig detrusor smooth muscle. *Am J Physiol Cell Physiol* **280**, C1090–C1096.
- ZhuGe R, Sims SM, Tuft RA, Fogarty KE & Walsh JV Jr (1998). Ca^{2+} sparks activate K^+ and Cl^- channels, resulting in spontaneous transient currents in guinea-pig tracheal myocytes. *J Physiol* **513**, 711–718.

Acknowledgements

The authors wish to acknowledge support from grant number 064212 from the Wellcome Trust, NIH RO1 DK68565 and the MRC. G.P.S. is in receipt of a Research Fellowship awarded by the Health Research Board, Ireland.

Ca²⁺ signalling in urethral interstitial cells of Cajal

Gerard P. Sergeant, M. A. Hollywood, N. G. McHale and K. D. Thornbury

J. Physiol. 2006;576;715-720; originally published online Aug 17, 2006;

DOI: 10.1113/jphysiol.2006.115956

This information is current as of November 13, 2007

Updated Information & Services	including high-resolution figures, can be found at: http://jp.physoc.org/cgi/content/full/576/3/715
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Review articles http://jp.physoc.org/cgi/collection/review_articles
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://jp.physoc.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://jp.physoc.org/misc/reprints.shtml